


【 Connecting institutions 】	【 Comments 】
<p>Peking Union Medical College Hospital (China), Fudan University Zhongshan Hospital (China), Tianjin Medical University (China), Fukuoka University Chikushi Hospital (Japan), Kyushu University Hospital (Japan), Shandong Province Hospital (China), Juntendo University (Japan), Beijing Hospital (China), Beijing Friendship Hospital, Capital Medical University (China)</p>	<p>In this conference, we discussed four EGC cases with special types and endoscopic manifestations. Gastric adenocarcinoma of fundic-gland mucosa type (GA-FGM) is sometimes difficult to identify under endoscopy. Remedial surgery is required when GA-FGM combined with deep infiltration in the submucosa because of unsatisfied prognosis. Another topic we discussed is how to identify adenomas and HGIN under endoscopy and pathology.</p>
	
<p>Teleconference view at Peking Union Medical College Hospital</p>	<p>Teleconference view at Tianjin Medical University</p>
<p>Picture taken at : Kyushu University Hospital</p>	<p>Picture taken at : Kyushu University Hospital</p>
	
<p>A presented slide</p>	<p>A presented slide</p>
<p>Picture taken at : Kyushu University Hospital</p>	<p>Picture taken at : Kyushu University Hospital</p>
<p>Magnifying Endoscopy</p>  <p>color difference boundary between the lesion and the surrounding mucosa microsurface structure varies in size but in a regular manner slightly widened intervening part no obvious irregular microvascular pattern</p>	 <p>Takashi Yao</p>
<p>A presented slide</p>	<p>Dr. Yao makes a comment</p>
<p>Picture taken at : Kyushu University Hospital</p>	<p>Picture taken at : Kyushu University Hospital</p>