





#2019

2025.04.14

The 46th Peking Early Gastric Cancer Tele-conference (PUMCH)

【 Connecting institutions 】	【 Comments 】
Peking Union Medical College H(China), Xizang Autonomous Region People's H (China), Shenzhen U (China), Nanjing Drum Tower H (China), Cancer H, Chinese Academy of Medical Sciences (China), Beijing H (China), Juntendo U (Japan), Shandong Province H (China), Kyushu U H (Japan), Fukuoka U Chikushi H (Japan), Tianjin Medical U (China), Fudan U Zhongshan H (China), Tibetan H of Tibet Autonomous Region (China), Olympus Latin America (United States)	The focus of this meeting was on the assessment of the differentiation degree of early cancer in the cardia region using magnifying gastroscopy, the follow-up treatment plan of GA-FGM for non-curative resection, and the emphasis on improving the focusing ability of magnifying gastroscopy to determine the DL. For EGC patients with current Hp infection, Hp eradication treatment should be performed after ESD.
	
A presentation slide Picture taken at：Kyushu University Hospital	A presentation slide Picture taken at：Kyushu University Hospital
	
A presentation slide Picture taken at：Kyushu University Hospital	A presentation slide Picture taken at：Kyushu University Hospital
	
A presentation slide Picture taken at：Kyushu University Hospital	A presentation slide Picture taken at：Kyushu University Hospital