




#2037

2025.07.07

The 47th Peking Early Gastric Cancer Tele-conference (PUMCH)

【 Connecting institutions 】	【 Comments 】
Peking Union Medical College H (China), Shandong Province H (China), Kyushu University H (Japan), Beijing Friendship H, Capital Medical University (China), Juntendo University (Japan), Nanfang H, Southern Medical University (China), Beijing H (China), Fukuoka University Chikushi H (Japan), Jiamusi University First Affiliated H (China), Tianjin Medical University (China), Tibetan H of Tibet Autonomous Region (China)	This meeting discussed four cases of early gastric cancer. Regarding GA-FG with pigmentation and glandular cystic dilation, it is very difficult to determine the DL under endoscopy. As to a minute gastric cancer, only scar changes by biopsy were observed in magnifying gastroscopy, but cancer was found in ESD-resected specimen. We discussed the pathological types of protruding ECG and endoscopic manifestations of HP-infected metachronous carcinoma.
	
Monitor shows connected sites	A presentation slide
Picture taken at：Kyushu University Hospital	Picture taken at：Kyushu University Hospital
	
A presentation slide	A presentation slide
Picture taken at：Kyushu University Hospital	Picture taken at：Kyushu University Hospital
	
A presentation slide	A presentation slide
Picture taken at：Kyushu University Hospital	Picture taken at：Kyushu University Hospital