




#2069

2025.09.29

The 48th Peking Early Gastric Cancer Tele-conference (PUMCH)

【 Connecting institutions 】	【 Comments 】
Peking Union Medical College H (China), Kyushu U H (Japan), Shandong Province H (China), Cancer H, Chinese Academy of Medical Sciences (China), Juntendo U (Japan), Fukuoka U Chikushi H (Japan), Nanfang H, Southern Medical U (China), Tianjin Medical U (China), Xizang Autonomous Region People's H (China), Fujian Provincial H (China), Fudan U Zhongshan H (China), Fourth H of Hebei Medical U (China), South China H of Shenzhen U (China)	For mixed undifferentiated EGC larger than 2cm, surgical salvage surgery is still recommended after ESD resection. The lesions that occur in the background of autoimmune gastritis are difficult to distinguish endoscopically as NET or GA-FG, and can only be confirmed by pathology. When the differentiation degree of internal components in EGC is unevenly distributed, it is not possible to directly confirm the degree of differentiation through magnifying gastroscopy.
	
Monitor shows connected sites	A presentation slide
Picture taken at：Kyushu University Hospital	Picture taken at：Kyushu University Hospital
	
A presentation slide	A presentation slide
Picture taken at：Kyushu University Hospital	Picture taken at：Kyushu University Hospital
	
A presentation slide	A presentation slide
Picture taken at：Kyushu University Hospital	Picture taken at：Kyushu University Hospital